

US009990581B2

(12) United States Patent

Carlsgaard et al.

(54) INSULIN DOSAGE ASSESSMENT AND RECOMMENDATION SYSTEM

(75) Inventors: Peter B. Carlsgaard, Zionsville, IN

(US); Christen Rees, Indianapolis, IN (US); Robin S. Wagner, Indianapolis,

ÍN (US)

(73) Assignee: Roche Diabetes Care, Inc.,

Indianapolis, IN (US)

(*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 798 days.

(21) Appl. No.: 13/546,447

(22) Filed: Jul. 11, 2012

(65) **Prior Publication Data**

US 2014/0019396 A1 Jan. 16, 2014

(51) Int. Cl.

 G06N 5/02
 (2006.01)

 G06F 19/00
 (2018.01)

 A61M 5/172
 (2006.01)

(52) U.S. Cl.

12/2/00

(58) Field of Classification Search

None

See application file for complete search history.

(56) References Cited

U.S. PATENT DOCUMENTS

7,404,796 B2 7/2008 Ginsberg 2003/0216628 A1 11/2003 Bortz et al.

(10) Patent No.: US 9,990,581 B2 (45) Date of Patent: Jun. 5, 2018

OTHER PUBLICATIONS

Thorne, "Attitudes toward patient expertise in chronic illness," Intl J Nursing Studies, 37, 303-311, 2000.*

Hornsten, "Nurses' experiences of conflicting encounters in diabetes care," EDN Summer, 5(2), 64-69, 2008.*

DAFNE Study Group, "raining in flexible, intensive insulin management to enable dietary freedom in people with type 1 diabetes: dose adjustment for normal eating (DAFNE) randomised controlled trial," vol. 325, 6 pages, 2002.*

(Continued)

Primary Examiner — G. Steven Vanni (74) Attorney, Agent, or Firm — Harness Dickey

(57) ABSTRACT

A computer-implemented method is presented for recommending insulin dosage adjustments for a patient having diabetes. The method includes: identifying a plurality of bolus events from patient data; grouping bolus events having a recommended bolus dosage substantially equivalent to the amount of administered insulin into a first subset of bolus events; determining a bolus outcome for each of the bolus events in the first subset of bolus events, where the bolus outcome is expressed in relation to a target range of blood glucose values and is selected from a group including above the target range, in the target range and below the target range; determining whether one of the bolus outcomes is predominant amongst the bolus events in the first subset of bolus events; and generating a recommendation pertaining to insulin dosage for the patient in response to a determination that one of the bolus outcomes is predominant.

13 Claims, 8 Drawing Sheets

